

# The routine immunisation schedule

from Autumn 2019

Age due	Diseases protected against	Vaccine given and trade name	Usual site
Eight weeks old	Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b (Hib) and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa Thigh
	Pneumococcal (13 serotypes)	Pneumococcal conjugate vaccine (PCV)	Prevenar 13 Thigh
	Meningococcal group B (MenB)	MenB	Bexsero Left thigh
	Rotavirus gastroenteritis	Rotavirus	Rotarix By mouth
Twelve weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa Thigh
	Rotavirus	Rotavirus	Rotarix By mouth
Sixteen weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa Thigh
	Pneumococcal (13 serotypes)	PCV	Prevenar 13 Thigh
	MenB	MenB	Bexsero Left thigh
One year old (on or after the child's first birthday)	Hib and MenC	Hib/MenC	Menitorix Upper arm/thigh
	Pneumococcal	PCV	Prevenar 13 Upper arm/thigh
	Measles, mumps and rubella (German measles)	MMR	MMR VaxPRO <sup>2</sup> or Priorix Upper arm/thigh
	MenB	MenB booster	Bexsero Left thigh
Eligible paediatric age groups <sup>1</sup>	Influenza (each year from September)	Live attenuated influenza vaccine LAIV <sup>2, 3</sup>	Fluenz Tetra <sup>2, 3</sup> Both nostrils
Three years four months old or soon after	Diphtheria, tetanus, pertussis and polio	DTaP/IPV	Infanrix IPV or Repevax Upper arm
	Measles, mumps and rubella	MMR (check first dose given)	MMR VaxPRO <sup>2</sup> or Priorix Upper arm
Boys and girls aged twelve to thirteen years	Cancers caused by human papillomavirus (HPV) types 16 and 18 (and genital warts caused by types 6 and 11)	HPV (two doses 6-24 months apart)	Gardasil Upper arm
Fourteen years old (school year 9)	Tetanus, diphtheria and polio	Td/IPV (check MMR status)	Revaxis Upper arm
	Meningococcal groups A, C, W and Y disease	MenACWY	Nimenrix or Menveo Upper arm
65 years old	Pneumococcal (23 serotypes)	Pneumococcal Polysaccharide Vaccine (PPV)	Pneumococcal Polysaccharide Vaccine Upper arm
65 years of age and older	Influenza (each year from September)	Inactivated influenza vaccine	Multiple Upper arm
70 years old	Shingles	Shingles	Zostavax <sup>2</sup> Upper arm

1. See Green book chapter 19 or visit [www.gov.uk/government/publications/influenza-the-green-book-chapter-19](http://www.gov.uk/government/publications/influenza-the-green-book-chapter-19) or [www.nhs.uk/conditions/vaccinations/child-flu-vaccine/](http://www.nhs.uk/conditions/vaccinations/child-flu-vaccine/)

2. Contains porcine gelatine.

3. If LAIV (live attenuated influenza vaccine) is contraindicated and child is in a clinical risk group, use inactivated flu vaccine.